

VERSION 02/06/16

**Trends in critical load exceedances in the UK
Updated June 2016**

Report to Defra, prepared under Contract AQ0826

Jane Hall¹ & Ron Smith²

¹CEH, Environment Centre Wales, Bangor, Gwynedd, LL57 2UW

²CEH, Bush Estate, Penicuik, Midlothian, EH26 0QB

CONTENTS	PAGE
Executive Summary	
1. Introduction	1
1.1 Overview of UK critical loads	1
1.1.1 Acidity critical loads	2
1.1.2 Nutrient nitrogen critical loads	3
1.2 Overview of UK deposition data	3
1.3 Overview of the calculation of critical load exceedances	4
1.3.1 Critical load exceedance metrics	5
1.3.2 Critical load exceedance maps for all habitats combined	6
2. Trends in critical loads exceedance by country and habitat	8
2.1 Trends by country	9
2.1.1 Acidity results	9
2.1.2 Nutrient nitrogen results	12
2.2 Trends by habitat	15
2.2.1 Acidity results	15
2.2.2 Nutrient nitrogen results	15
3. Trends in exceedance of site-relevant critical loads (SRCL)	22
3.1 Overview of site-relevant critical loads	22
3.2 Overview of SRCL exceedance metrics	23
3.2.1 Acidity results	24
3.2.2 Nutrient nitrogen results	29
References	34

Executive Summary

Critical loads define the amount of acid or nitrogen deposition below which significant harmful effects do not occur to sensitive habitats. An “exceedance” is the amount of excess acid or nitrogen deposition above the critical load. This report presents the trends in critical load exceedances for UK broad habitats, based on deposition data covering the period from 1995 to 2013. Summary statistics are published to monitor progress in the areas at risk from air pollution over time, and are used for:

- Defra: Environmental Statistics – Key Facts
<https://www.gov.uk/government/publications/environment-statistics-key-facts>
- Welsh Government: Sustainable Development Indicators for Wales
<http://wales.gov.uk/topics/statistics/headlines/sustaindev/120829/?lang=en>
- Scottish Government: Key Scottish Environment Statistics
<http://www.scotland.gov.uk/Topics/Statistics/Browse/Environment/>
- UK Biodiversity Indicators in Your Pocket: JNCC; biodiversity indicator for assessing the pressures from air pollution
<http://jncc.defra.gov.uk/page-4233>

For acidity, the area of sensitive habitats in the UK with exceedance of critical loads has fallen by 28.5% since 1995, from 72.6% based on 3-year mean deposition data for 1995-97, to 44.1% based on mean deposition data for 2012-14. Over the same time period the Average Accumulated Exceedance has more than halved, from 0.78 to 0.28 keq ha⁻¹ year⁻¹.

For nutrient nitrogen, the reductions are smaller with a 12.8% decrease in the area exceeding nitrogen critical loads, from 75% in 1995-97 to 62.2% using deposition data for 2012-14. The Average Accumulated Exceedance for nutrient nitrogen has declined by one third, from 9.5 kg N ha⁻¹ year⁻¹ to 6.0 kg N ha⁻¹ year⁻¹ over the same time period.

Additionally this report now includes trends in exceedances of critical loads applied to the features of designated sites (Special Areas of Conservation: SACs; Special Protection Areas: SPAs; Sites of Special Scientific Interest: SSSIs). For SACs the percentage of sites with exceedance of acidity critical loads for any feature has decreased by 15% between 1995 and 2014, and by 24% for SPAs and 16% for SSSIs. Over the same time period the maximum Average Accumulated Exceedance has fallen by more than 50%. For nutrient nitrogen the reductions are smaller, between 5 and 10% reduction in the percentage of sites with exceedance for any feature, and a 35% reduction in the maximum Average Accumulated Exceedance.

1. Introduction

This report presents the trends in the areas of sensitive habitats at risk from the adverse impacts of excess acid and/or nitrogen deposition. The metrics are based on the exceedance of critical loads for acidification and eutrophication. This section provides a brief overview of UK critical loads and deposition data and the calculation of exceedances; further details can be found in Hall et al (2014). The trends in critical loads exceedances are presented and discussed in Section 2. The application of “site-relevant critical loads” (SRCL) for designated sites, and trends in their exceedances are described in Section 3.

1.1 Overview of UK critical loads

Critical loads are thresholds for effects from atmospheric deposition and defined as “a quantitative estimate of the exposure to one or more pollutants below which significant harmful effects on specified sensitive elements of the environment do not occur according to present knowledge” (Nilsson & Grennfelt, 1988).

The methods used to calculate and map UK critical loads are described in detail in Hall et al (2015). Critical loads are calculated and mapped for UK habitats sensitive to acidification and/or eutrophication (Table 1.1).

Table 1.1: Habitat distributions mapped for acidity and for nutrient nitrogen critical loads

Habitat	EUNIS habitat class(es) assigned ¹	Mapped for acidity	Mapped for nutrient nitrogen
Acid grassland (wet & dry)	E1.7 & E3.52	Yes	Yes
Calcareous grassland	E1.26	Yes	Yes
Dwarf shrub heath (wet & dry)	F4.11 & F4.2	Yes	Yes
Montane	E4.2	Yes	Yes
Bog	D1	Yes	Yes
Managed coniferous woodland	G3	Yes	Yes
Managed broadleaved woodland	G1	Yes	Yes
Beech woodland (unmanaged)	G1.6	Yes (mapped together)	Yes
Acidophilous oak woodland (unmanaged)	G1.8		Yes
Scots Pine woodland (unmanaged)	G3.4		Yes
Other unmanaged woodland	G4		Yes
Freshwaters ²	C1 & C2	Yes	No
Dune grassland	B1.4	No	Yes
Saltmarsh	A2.53/54/55	No	Yes

¹EUNIS class closest to broad habitat and critical loads habitat; class used for assigning empirical nutrient nitrogen critical loads and for classifying UK critical loads data for submission to the CCE.

²Critical loads are calculated for 1752 freshwater sites across the UK (see Section 1.1.1 below); habitat areas are based on the catchment areas of these sites.

Published correspondence tables (available from: <http://jncc.defra.gov.uk/page-1425>) are used to relate broad habitats to the European Nature Information System (EUNIS: Davies & Moss, 2002) hierarchical habitat classification scheme, developed for pan-European applications.

Habitat distribution maps are based on the CEH Land Cover Map 2000 (LCM2000: Fuller et al, 2002(a)(b)) and additional data sets such as species distribution data and altitude. Habitat areas, used for assessing the areas of habitats at risk from acidification and or eutrophication, are based on the LCM2000 data. It should be noted that the habitat distribution maps and areas used for UK critical loads (acidity, nitrogen) research (a) only include areas where data exist for the calculation or derivation of critical loads; (b) may differ from other national habitat distribution maps or estimates of habitat areas. This may also result in a difference in the total habitat areas mapped for acidity and for nutrient nitrogen critical loads.

1.1.1 Acidity critical loads

Two methods are used in the UK for calculating acidity critical loads for terrestrial habitats: the empirical approach is used to provide estimates for non-woodland habitats and a simple mass balance equation used for woodland habitats.

An empirical approach is used to define acidity critical loads for UK soils; critical loads are assigned to each 1km grid square of the UK based on the amount of acid deposition that could be neutralised by the base cations produced by mineral weathering of the dominant soil type in the grid square. This approach is applied to mineral and organo-mineral soils (Hornung et al, 1995) but is inappropriate for peat soils because of the absence of inputs of alkalinity from mineral weathering (Smith et al, 1992; Gammack et al, 1995). Critical loads of acidity for peat soils are set to the value corresponding to the amount of acid deposition that would give rise to an effective rain pH value of 4.4 (Calver, 2003; Calver et al, 2004; Skiba & Cresser, 1989); this pH reflects the buffering effects of organic acids upon peat drainage water pH. This method is applicable to upland and lowland acid peat soils, but not to peats in lowland arable fen areas that are less sensitive to acidification, where a higher critical load is set than would be applied to acid peats (Hall et al, 2015).

The acidity critical loads for soils as outlined above are used to set the acidity critical loads to protect the soils on which non-woodland habitats occur. In addition, they are used, with additional habitat-specific data, in deriving the acidity critical load input parameters for the “Critical Loads Function” (Section 2).

For woodland habitats a simple mass balance (SMB) equation, based on balancing the acidic inputs to and outputs from the ecosystem, is used to derive a critical load that ensures a specified critical chemical limit is not exceeded (Sverdrup et al, 1990; Sverdrup & De Vries, 1994). In the UK the SMB equation is parameterised using different chemical criteria for woodlands on mineral or organo-mineral soils, and woodlands on peat soils (Hall et al, 2015). Critical loads are calculated for both managed (productive) and unmanaged woodlands in order to protect the long-term ecosystem function of the woodland habitats; this also aims to protect the land under managed conifer forest for possible future non-forest use and reversion to semi-natural land uses. These critical loads are also used with additional habitat-specific data to derive the acidity critical load input parameters for the “Critical Loads Function” (Section 2) for woodland habitats.

Acidity critical loads for freshwaters are calculated using the catchment-based First-Order Acidity Balance (FAB: Henriksen & Posch, 2001) model. FAB is currently applied to 1752 sites across the UK, comprising a mixture of mainly upland, lakes, reservoirs and first-order streams (ie, streams that

feed into other larger streams, but do not have any other streams draining into them). The critical load calculations are based on the most recent, best available estimate of annual mean water chemistry data.

1.1.2 Nutrient nitrogen critical loads

Empirical and mass balance methods also exist for calculating critical loads for eutrophication (ie, an excess of nitrogen as a nutrient). The empirical critical loads are based on experimental or field evidence of thresholds for changes in species composition, plant vitality or soil processes. The empirical approach is suited to semi-natural communities for which the long-term protection of biodiversity and/or ecosystem function is the key concern. In the UK the empirical approach is applied to natural and semi-natural habitats, including unmanaged (non-productive) woodland, based on critical load values agreed at international workshops (Bobbink & Hettelingh, 2011; Hall et al, 2015).

In the mass balance approach the long-term inputs and outputs of nitrogen from the ecosystem are calculated, with the critical load being exceeded when any excess nitrogen input is calculated to lead to an exceedance of a specified critical rate of nitrogen leaching. This approach is suited to managed ecosystems of low biodiversity, in which the inputs and outputs can be quantified with some confidence and in which the key concern is nitrate leaching. In the UK, this approach is applied to managed (productive) woodlands to ensure that long-term ecosystem function (eg, soils, soil biological resources, trees, linked aquatic systems) is protected.

1.2 Overview of UK deposition data

The sulphur, nitrogen and base cation deposition data used in the UK calculations of critical loads and their exceedances are based on the “Concentration Based Estimated Deposition” (CBED) methodology (RoTAP, 2012). Site based measurements of air concentrations of sulphur and nitrogen gases are interpolated to generate 5km maps of concentrations for the UK. Ion concentrations in precipitation (from the UK Eutrophying and Acidifying Pollutants (UKEAP) network) are combined with the Met Office annual precipitation map to generate maps of wet deposition. The wet deposition values include (a) direct deposition of cloud droplets to vegetation (known as “occult” deposition); (b) an orographic enhancement to take account of the “seeder-feeder” effect in upland regions (Fowler et al, 1988). Gas and particulate concentration maps are combined with spatially distributed estimates of vegetation-specific deposition velocities (Smith et al, 2000) to generate dry deposition. Combining these data sets produces 5km maps of total (wet + cloud + dry) deposition of sulphur (non-marine), oxidised nitrogen and reduced nitrogen; two different sets of deposition values are used in critical load and exceedance applications: (i) assumes grassland or moorland vegetation everywhere; (ii) assumes forest everywhere, based on the different deposition velocities to different land cover types.

Significant inter-annual variations in deposition can occur due to the natural variability in annual precipitation (which influences wet deposition) as well as the general circulation of air which can increase or decrease the amount of polluted air imported from the European continent. The CBED deposition data used to calculate critical load exceedances is therefore averaged over a three-year period; this has been demonstrated to be a suitable time period to smooth out inter-annual variations in deposition.

As critical loads for terrestrial habitats are mapped on a 1km grid, for exceedance calculations deposition is assumed to be constant for all 1 km squares within each 5km grid square. For freshwater exceedance calculations catchment-weighted mean sulphur and nitrogen deposition values are calculated by overlaying the catchment boundary and land cover information (moorland vs forest) onto the 5km deposition maps.

1.3 Overview of the calculation of critical load exceedances

Critical load exceedances are the amount of excess deposition above the critical load; for nutrient nitrogen the calculation is simply total nitrogen deposition (derived from nitrogen oxides and ammonia) minus the critical load. For acidification, deposition of both sulphur and nitrogen compounds can contribute to the exceedance of critical loads. The Critical Load Function, developed under the UNECE CLRTAP (Posch *et al.*, 1999; Posch & Hettelingh, 1997; Posch *et al.*, 1995; Hettelingh *et al.*, 1995), defines combinations of sulphur and nitrogen deposition that will not cause harmful effects. In its simplest form, an acidity critical load can be defined graphically by a 45 degree diagonal line on a sulphur-nitrogen deposition plot (Figure 1.1a). The line intercepts the x-axis (representing nitrogen deposition) and y-axis (representing sulphur deposition) at chemically equivalent points, each representing the nitrogen or sulphur deposition equal to the critical load for acidity. Each point along the diagonal line represents the critical load in terms of some combination of sulphur and nitrogen deposition.

To allow for the long-term nitrogen removal processes by the soil and through harvesting of vegetation, the simple diagonal line is shifted along the nitrogen axis to increase the nitrogen values across the entire CLF (Figure 1.1b). More nitrogen can then be deposited before the acidity critical load is exceeded. There are no similar removal processes that need to be considered for sulphur.

The intercepts of the CLF on the sulphur and nitrogen axes (Figure 1.1c) define the following terms:

- The “maximum critical load of sulphur” (CL_{maxS}): the critical load for acidity expressed in terms of sulphur only, ie, when nitrogen deposition is zero.
- The “maximum critical load of nitrogen” (CL_{maxN}): the critical load for acidity expressed in terms of nitrogen only (when sulphur deposition is zero).
- The “minimum critical load of nitrogen” (CL_{minN}): the long-term nitrogen removal processes in the soil (eg, nitrogen uptake and immobilisation) and harvesting of vegetation.

These critical loads are calculated from the acidity critical loads described in Section 1.1 and additional soil-specific or habitat-specific data.




Figure 1.1: Development of the CLF: (a) acidity critical load defined by equal amounts of sulphur and nitrogen deposition; (b) shifting the acidity critical load diagonal line to allow for nitrogen removal

processes; (c) the 3 nodes of the CLF: CLmaxS, CLminN, CLmaxN. The area shown in grey represents the combinations of sulphur and nitrogen deposition that are below the critical load (ie, critical load is not exceeded).

Exceedances are calculated by comparing the values of CLmaxS, CLminN and CLmaxN to the values of sulphur and nitrogen (oxidised + reduced) deposition. The actual calculation depends on where the deposition falls in relation to these critical load values; the CLF is divided into five different regions for this purpose (Figure 1.2). The exceedance is defined by the sum of sulphur and nitrogen deposition as shown by the red arrows in Figure 1.2 (ie, not the length of the diagonal line); this is referred to as the “shortest distance” exceedance. Further details on the calculations are given in Hall et al (2015).




Figure 1.2: Example of S and N deposition reductions required depending on the region of the CLF. Deposition that falls in region 5 is below the critical load (ie, critical loads not exceeded).

1.3.1 Critical load exceedance metrics

Critical load exceedances are calculated for each 1km square of the distributions of each terrestrial habitat, and for each catchment for freshwaters. The results are then summarised by habitat and country using the following exceedance metrics:

(i) Area of habitat exceeded

For terrestrial habitats the area values are based on the LCM2000 data; if the critical load for any individual habitat is exceeded, the exceeded area is set to the habitat area within the 1km square for that particular habitat. For freshwater habitats, if the FAB critical load is exceeded, the whole catchment is assumed to be exceeded and the exceeded area set to the catchment area. The total exceeded areas for individual habitats are summarised by country.

(ii) Percentage area of habitat exceeded

This is calculated from the exceeded areas derived in (i) and the total area of each habitat mapped in each country (Section 1.1). While this is a useful metric, it has its limitations, for example, when comparing exceedance results from one year to another (or one deposition scenario to another), there may very small (or no) changes in the percentage area of habitat exceeded. This is because the magnitude of the exceedance may have reduced, but the area

exceeding the critical load remains the same; the area exceeded will only reduce when the critical load is no longer exceeded.

(iii) Accumulated Exceedance (AE)

AE takes account of both the magnitude of exceedance and the habitat area exceeded:

$$AE \text{ (keq year}^{-1}\text{)} = \text{exceedance (keq ha}^{-1} \text{ year}^{-1}\text{)} * \text{exceeded area (ha)}$$

AE is calculated for each 1km square for each habitat and then summarised by habitat and country. AE is set to zero where critical loads are not exceeded. This metric can be useful for comparing results for different years or scenarios, but because the results are expressed in keq year⁻¹ they tend to be very large numbers and not intuitive to understand. It should also be noted that the same AE can arise from a large exceedance and small exceeded area, or a small exceedance and a large area.

(iv) Average Accumulated Exceedance (AAE)

AAE averages the AE across the entire sensitive habitat area:

$$AAE \text{ (keq ha}^{-1} \text{ year}^{-1}\text{)} = AE \text{ (keq year}^{-1}\text{)} / \text{total habitat area (ha)}$$

This metric provides an exceedance value averaged across the whole habitat area. In the summary statistics presented (Section 2) it is based on the AE for the habitat (by country) divided by the total habitat area (by country). AAE is set to zero where critical loads are not exceeded. This metric provides a more intuitive value for comparing the exceedance results for different years or scenarios, and gives an indication of the reduction in the magnitude of exceedance even if there is no change in the percentage area of habitat exceeded.

1.3.2 Critical load exceedance maps for all habitats combined


Critical load exceedances are calculated by habitat; exceedance maps can be generated for individual habitats or for all terrestrial habitats combined. The exceedance data for freshwaters are not incorporated into these combination maps because the data are catchment-based rather than for 1km squares and as such may overlap with other habitat data. This section focuses on maps of AAE for all terrestrial habitats combined (Figure 1.3); other maps are presented and discussed in Hall et al (2014). Maps of AAE provide a good representation of the summary critical load exceedance statistics since they are based on all the critical load values for all habitats and habitat-specific deposition. The AAE for each 1km square is calculated as:

$$AAE = \sum(AE \text{ for all habitats}) / \sum(\text{area for all habitats})$$

AE (and AAE) is set to zero where the critical loads are not exceeded.

The latest AAE maps for acidity and nutrient nitrogen (Figure 1.3) clearly show the lower exceedances in Scotland compared to other regions of the UK. High exceedances of acidity critical loads are focused in upland areas of central and north western England, as well as smaller areas in eastern England and the far south-west, as well as parts of Wales and southern Scotland and Northern Ireland. High exceedances of nutrient nitrogen critical loads are widespread across England, Wales and Northern Ireland and parts of southern and eastern Scotland, with many areas having exceedances above 14 kg N ha⁻¹ year⁻¹ (1 keq ha⁻¹ year⁻¹).

(a) Acidity



(b) Nutrient nitrogen




Figure 1.3: Average Accumulated Exceedance (AAE) of critical loads by CBED deposition for 2012-14. Although the legends for the two maps are given in different units the class intervals are equivalent (ie, 7 kg N ha⁻¹ year⁻¹ is equivalent to 0.5 keq ha⁻¹ year⁻¹).

2. Trends in critical loads exceedance by habitat and country

Acidity and nutrient nitrogen exceedances by habitat and country are updated annually using the latest 3-year rolling mean CBED deposition data. The summary statistics as described in Section 1.3.1 are made available to Defra and the Devolved Administrations and JNCC; from these they have used the trends in the percentage area of habitats exceeded for the following:

- Defra: Environmental Statistics – Key Facts
<https://www.gov.uk/government/publications/environment-statistics-key-facts>
- Welsh Government: Sustainable Development Indicators for Wales
<http://wales.gov.uk/topics/statistics/headlines/sustaindev/120829/?lang=en>
- Scottish Government: Key Scottish Environment Statistics
<http://www.scotland.gov.uk/Topics/Statistics/Browse/Environment/>
- JNCC: biodiversity indicator for assessing the pressures from air pollution
<http://jncc.defra.gov.uk/page-4233>

The data used for the trends analysis are summarised in Box 1; there are a few inconsistencies between years due to changes in methods used to derive deposition estimates, and some minor alterations to the acidity critical loads. This information should be taken into account when interpreting the trends results.

Box 1:

Data used for critical loads trends analysis

Critical loads data

Acidity: data as summarised in Section 1.1.1 of this report were used for all years except results prior to 2004-2006 where: (a) the acidity critical loads for the bog habitat were based on the dominant soil in each 1x1km grid square; later results use critical loads data that assume all areas of bog habitat occur on peat soils; (b) freshwater exceedances were based on catchment-weighted grid-average deposition; the later results are based on catchment-weighted ecosystem-specific deposition. Note that the freshwater results are based on critical loads for 1752 lake or stream sites across the UK, and therefore do not represent all waters in the UK.

Nutrient nitrogen: data as summarised in Section 1.1.2 of this report.

Deposition data

All results based on 5x5 km resolution “concentration based estimated deposition” (CBED) values averaged over a three year period. All data are based on a consistent methodology except:

- (a) Deposition data prior to 2001-2003 exclude nitric acid as the monitoring network for this pollutant was not in operation prior to this time.
- (b) Deposition data prior to 2002-2004 excludes aerosol deposition of NH_4 , NO_3 , SO_4 .
- (c) Data for 2004-06 onwards updated in February 2015 to correct for over-estimate of nitric acid deposition.

CBED moorland values are applied to non-woodland terrestrial habitats, and CBED woodland values are applied to woodland habitats.

Habitat area data

These are based on the habitat distribution maps generated for UK critical loads research (see Section 1.1 of this report). There was a small reduction in the area mapped for acidity for the bog habitat as a result of the change to the critical loads in 2008; results using the updated habitat area apply to all results from 2004-06 onwards.

The trends results are shown as both tables and simple plots; it is worth noting that while the percentage area exceeded for some habitats may not alter from one year to another, the AE values fluctuate reflecting changes in the national deposition data.

2.1 Trends by country

Table 2.1 shows the total land area by country and the area of habitats sensitive to acidification and eutrophication to which critical loads have been applied; 31% of the UK land area has habitats mapped for acidity critical loads, and 29% for nutrient nitrogen. *Note: throughout this report the summary exceedance statistics of the percentage area exceeded are percentages of the habitat areas mapped as sensitive to acidification/eutrophication (ie, not % land area).*

Table 2.1: Total land area and habitat areas mapped for critical loads by country

Country	Land area (km ²) [#]	Habitat areas mapped for acidity (km ²)	Area mapped for acidity as % of country	Habitat areas mapped for nutrient nitrogen (km ²)	Area mapped for nutrient nitrogen as % of country
England	132938	18635	14	19522	15
Wales	21225	7798	37	6837	32
Scotland	80239	48083	60	43200	54
NI	14130	3541	25	3467	25
UK	248532	78051	31	73027	29

[#] The UK and its countries: facts and figures. Office for National Statistics:

<http://webarchive.nationalarchives.gov.uk/20160105160709/http://www.ons.gov.uk/ons/guide-method/geography/beginner-s-guide/administrative/the-countries-of-the-uk/index.html>

2.1.1 Acidity results

The results for acidity (Table 2.2, Figure 2.1) show that the total area of habitats exceeding critical loads in the UK has declined from 72.6% in 1995-97 to 44.1% in 2012-14. However, the area exceeded varies between countries (Table 2.2, Figure 2.2), due to (a) geographic location of different sensitive habitats across the country (see Section 2.2); (b) the range in critical load values across the country – lower critical loads are mainly found in the uplands in the north and west in the UK; (c) higher wet deposition (and therefore higher total deposition) in the uplands or wetter regions of the country. The percentage area of habitats exceeded is lowest in Scotland in all years; however as shown in Table 2.1 61.1% of Scotland has habitats mapped for acidity critical loads, and that means the actual areas exceeded are larger than in the other countries (eg, 14856 km² exceeded by 2012-14 deposition). Although only 14.3% of England has habitats mapped for acidity critical loads, 61.6% of their area is exceeded for 2012-14, equivalent to 11483 km². The magnitude of exceedance across the UK, expressed as AAE (Table 2.3, Figure 2.1), has more than halved from 0.78 keq ha⁻¹ year⁻¹ in 1995-97 to 0.28 keq ha⁻¹ year⁻¹ in 2012-14. The data show the largest reductions in the exceedances were in the late 1990s; changes since then have been smaller and fluctuated from one year to another, but continuing the general downward trend. Note that the acidity critical loads for calcareous grassland are not exceeded in any year (Table 2.3).

Table 2.2: Acidity: Percentage area of habitats by country and deposition dataset year where acidity critical loads are exceeded

Year	Percentage habitat area exceeded by country:				
	England	Wales	Scotland	NI	UK
1995-1997	75.8	90.0	68.2	76.8	72.6
1998-2000	71.6	83.1	52.6	67.2	60.8
1999-2001	71.9	83.0	51.6	66.8	60.3
2001-2003	72.3	82.4	43.0	67.4	55.0
2002-2004	72.3	82.3	44.8	69.2	56.2
2003-2005	71.8	83.2	44.5	67.1	55.9
2004-2006	66.8	81.2	48.0	68.1	56.7
2005-2007	66.1	81.0	46.1	68.5	55.4
2006-2008	64.3	79.2	40.7	68.6	51.4
2007-2009	63.6	77.4	32.9	69.4	46.3
2008-2010	63.2	74.9	31.5	69.6	45.2
2009-2011	63.8	74.5	33.9	71.0	46.8
2010-2012	62.8	74.2	32.2	67.8	45.3
2011-2013	62.1	74.4	31.0	69.4	44.5
2012-2014	61.6	75.3	30.9	63.4	44.1
Reduction in % area exceeded 1995-2014	14.2	14.7	37.3	13.4	28.5

Table 2.3: Acidity: Average Accumulated Exceedance (AAE in keq ha⁻¹ year⁻¹) by country and deposition dataset year

Year	AAE (keq ha ⁻¹ year ⁻¹) by country:				
	England	Wales	Scotland	NI	UK
1995-1997	1.33	1.36	0.47	0.80	0.78
1998-2000	1.00	0.84	0.28	0.46	0.51
1999-2001	0.98	0.82	0.27	0.46	0.50
2001-2003	1.04	0.82	0.23	0.51	0.50
2002-2004	0.94	0.79	0.24	0.46	0.48
2003-2005	0.93	0.84	0.24	0.42	0.47
2004-2006	0.77	0.74	0.24	0.42	0.43
2005-2007	0.74	0.73	0.21	0.45	0.40
2006-2008	0.68	0.61	0.17	0.44	0.35
2007-2009	0.62	0.54	0.12	0.45	0.3
2008-2010	0.59	0.49	0.12	0.47	0.29
2009-2011	0.62	0.48	0.15	0.53	0.31
2010-2012	0.6	0.47	0.14	0.46	0.3
2011-2013	0.59	0.47	0.13	0.46	0.29
2012-2014	0.56	0.51	0.13	0.35	0.28
Reduction in AAE 1995-2013	0.77	0.85	0.34	0.45	0.50




Figure 2.1: Acidity: Percentage area of acid-sensitive habitats with exceedance of acidity critical loads in the UK by year, and AAE in $\text{keq ha}^{-1} \text{ year}^{-1}$.




Figure 2.2: Acidity: Percentage area of acid-sensitive habitats with exceedance of acidity critical loads, by country and year, and AAE in $\text{keq ha}^{-1} \text{ year}^{-1}$.

2.1.2 Nutrient nitrogen results

The results for nutrient nitrogen (Table 2.4 and Figure 2.3) show a decline in the percentage area of habitats exceeded in the UK, from 75% in 1995-97 to 62.2% in 2010-12. The results for England and Wales remained above, or close to, 90% exceeded over the same time period (Table 2.4, Figure 2.4). Scotland shows the smallest percentage habitat area exceeded of all countries, but the area exceeded (17579 km² for 2012-14) is similar to the area exceeded in England (18755 km² in 2010-12). The results reflect the smaller reductions in nitrogen deposition over the last two decades compared to the reductions in sulphur deposition (which helped reduce the exceedances of acidity critical loads). However, the magnitude of the exceedance (expressed as AAE) across the UK has reduced by one-third, from 9.5 kg N ha⁻¹ year⁻¹ in 1995-97 to 6.0 kg N ha⁻¹ year⁻¹ in 2012-14 (Table 2.5, Figure 2.3). The AAE varies from one region to another with the lowest values in Scotland and the highest in England (Table 2.5, Figure 2.4).

Table 2.4: Nutrient nitrogen: Percentage area of habitats by country and deposition dataset year where nutrient nitrogen critical loads are exceeded

Year	Percentage habitat area exceeded by country:				
	England	Wales	Scotland	NI	UK
1995-1997	98.3	98.0	59.4	92.6	75.0
1998-2000	97.6	92.5	48.9	80.0	67.5
1999-2001	97.7	91.1	50.9	82.5	68.7
2001-2003	97.8	93.5	47.7	85.4	67.1
2002-2004	97.6	93.3	50.2	86.3	68.6
2003-2005	97.5	94.1	50.6	83.8	68.8
2004-2006	96.7	93.2	52.9	84.8	69.9
2005-2007	96.5	93.6	53.6	86.4	70.4
2006-2008	96.1	92.9	49.0	86.8	67.5
2007-2009	96.4	91.7	41.8	88.7	63.3
2008-2010	96.5	89.7	40.7	89.7	62.6
2009-2011	97.0	89.8	44.5	91.4	65.0
2010-2012	96.5	89.6	41.4	88.5	62.9
2011-2013	96.0	90.3	40.7	89.9	62.5
2010-2014	96.1	90.9	40.7	83.0	62.2
Reduction in % area exceeded 1995-2013	2.2	7.1	18.7	9.6	12.8

Table 2.5: Nutrient nitrogen: Average Accumulated Exceedance (AAE in kg N ha⁻¹ year⁻¹) by country and deposition dataset year

Year	AAE (kg N ha ⁻¹ year ⁻¹) by country:				
	England	Wales	Scotland	NI	UK
1995-1997	19.0	15.8	4.1	10.6	9.5
1998-2000	16.8	10.3	2.7	6.5	7.4
1999-2001	17.4	10.6	2.9	6.8	7.7
2001-2003	19.7	12.2	3.1	8.9	8.7
2002-2004	18.0	12.2	3.3	8.7	8.3
2003-2005	18.2	13.2	3.3	8.3	8.4
2004-2006	14.9	11.4	3.1	7.9	7.2
2005-2007	14.9	11.4	2.9	8.8	7.2
2006-2008	14.1	9.9	2.5	8.8	6.6
2007-2009	13.8	9.5	2.1	9.4	6.3
2008-2010	13.9	9.2	2.2	9.8	6.3
2009-2011	14.6	9.2	2.6	10.9	6.8
2010-2012	13.8	8.8	2.4	9.6	6.4
2011-2013	13.3	8.9	2.3	9.5	6.2
2012-2014	12.7	9.1	2.3	7.6	6.0
Reduction in AAE 1995-2013	6.3	6.7	1.8	3.0	3.5




Figure 2.3: Nutrient nitrogen: Percentage area of nitrogen-sensitive habitats with exceedance of nitrogen critical loads in the UK by year, and AAE in kg N ha⁻¹ year⁻¹.




Figure 2.4: Nutrient nitrogen: Percentage area of nitrogen-sensitive habitats with exceedance of nitrogen critical loads, by country and year, and AAE in kg N ha⁻¹ year⁻¹.

2.2 Trends by habitat

Exceedances are summarised by habitat and country. This section focuses on the results by habitat for the UK; habitat results for individual countries are supplied to Defra and the devolved administrations.

2.2.1 Acidity results

As mentioned in Section 2.1 there is no exceedance of the acidity critical loads for calcareous grassland and this habitat is therefore excluded from Tables 2.6 and 2.7 and Figures 2.5 and 2.6. The habitats with the highest percentage area exceeded are acid grassland, montane, bog and managed woodlands (Table 2.6, Figure 2.5); these habitats also have the highest AAE values (Table 2.7, Figure 2.5). Of the habitats mapped for acidity, dwarf shrub heath is the habitat with the largest cover across the UK (10.1%); the largest decrease (42.4%) in the area exceeded is also seen for this habitat from 70.3% in 1995-97 to 27.9% in 2012-14. The largest reductions in AAE over the same timescale are for woodland, acid grassland and montane habitats (Table 2.7).

2.2.2 Nutrient nitrogen results

There are six habitats with more than 80% of their area exceeded for nitrogen in all years (Table 2.8, Figure 2.6): calcareous grasslands and woodlands (beech, oak, managed conifer and broadleaf and other unmanaged woodland). The largest reduction (44.1%) in the area exceeded is for dune grassland from 70.6% in 1995-97 to 26.5% in 2012-14. Another coastal habitat, saltmarsh, has virtually no exceedance in any year, due to a combination of its high critical load and the lower deposition in coastal areas. AAE is generally highest for the woodland habitats (Table 2.9, Figure 2.6), with the exception of Scots Pine, which is only found in Scotland where the magnitude of exceedance is generally lower due to the lower deposition in this region. The beech woodland is virtually 100% exceeded in all years, but the AAE has decreased from 22.7 kg N ha⁻¹ year⁻¹ in 1995-97 to 12.5 kg N ha⁻¹ year⁻¹ in 2012-14.

Table 2.6: Acidity: Percentage area of habitats where acidity critical loads are exceeded in the UK by deposition dataset year.

Year	Percentage habitat area with exceedance of acidity critical loads:								
	Acid grassland	Dwarf shrub heath	Bog	Montane	Coniferous woodland (managed)	Broadleaved woodland (managed)	Unmanaged woodland	Freshwaters	All habitats
1995-1997	92.0	70.3	88.0	95.8	79.4	75.8	69.5	29.9	72.6
1998-2000	84.9	49.5	78.8	91.3	69.9	68.4	57.2	24.2	60.8
1999-2001	84.7	47.9	76.1	93.4	70.2	69.1	58.2	23.9	60.3
2001-2003	79.5	40.7	61.6	82.5	66.5	69.6	58.5	21.9	55.0
2002-2004	80.3	42.3	60.1	89.4	68.4	70.8	60.1	21.3	56.2
2003-2005	80.4	41.5	59.3	92.9	68.2	70.1	59.1	21.7	55.9
2004-2006	82.5	45.1	71.7	96.3	64.2	61.5	48.3	21.7	56.7
2005-2007	81.9	41.5	76.4	94.4	63.6	60.5	46.7	21.3	55.4
2006-2008	78.9	35.4	73.3	85.6	60.5	57.2	43.4	20.6	51.4
2007-2009	73.9	28.5	63.7	71.4	57.1	55.9	42.1	19.0	46.3
2008-2010	72.2	28.1	57.9	70.1	55.6	55.6	42.0	18.5	45.2
2009-2011	74.8	30.6	54.9	71.6	58.0	57.0	43.3	18.9	46.8
2010-2012	73.0	29.3	54.4	65.3	56.7	55.0	41.5	19.0	45.3
2011-2013	73.4	28.5	50.1	62.3	56.2	53.5	40.5	18.8	44.5
2012-2014	71.8	27.9	56.7	60.4	55.7	52.4	38.9	18.9	44.1
Reduction in % area exceeded 1995-2013	20.2	42.4	31.3	35.4	23.7	23.4	30.7	11.0	28.4

Table 2.7: Acidity: AAE (in keq ha⁻¹ year⁻¹) by habitat for the UK by deposition dataset year.

Year	AAE (keq ha ⁻¹ year ⁻¹) by habitat:								
	Acid grassland	Dwarf shrub heath	Bog	Montane	Coniferous woodland (managed)	Broadleaved woodland (managed)	Unmanaged woodland	Freshwaters	All habitats
1995-1997	1.15	0.47	0.76	0.81	1.13	1.20	0.87	0.36	0.78
1998-2000	0.80	0.28	0.53	0.57	0.68	0.88	0.58	0.23	0.51
1999-2001	0.77	0.26	0.50	0.59	0.68	0.90	0.61	0.21	0.50
2001-2003	0.70	0.24	0.46	0.60	0.72	1.01	0.68	0.18	0.50
2002-2004	0.67	0.22	0.41	0.64	0.74	0.94	0.65	0.17	0.48
2003-2005	0.68	0.21	0.39	0.62	0.73	0.94	0.65	0.17	0.47
2004-2006	0.68	0.22	0.44	0.66	0.58	0.66	0.44	0.17	0.43
2005-2007	0.64	0.19	0.45	0.53	0.56	0.65	0.43	0.16	0.40
2006-2008	0.57	0.16	0.42	0.39	0.49	0.56	0.36	0.13	0.35
2007-2009	0.49	0.12	0.34	0.28	0.43	0.53	0.34	0.12	0.30
2008-2010	0.47	0.12	0.33	0.28	0.42	0.52	0.34	0.11	0.29
2009-2011	0.51	0.14	0.35	0.31	0.46	0.56	0.36	0.12	0.31
2010-2012	0.50	0.13	0.35	0.26	0.43	0.51	0.32	0.12	0.30
2011-2013	0.51	0.13	0.34	0.25	0.42	0.47	0.30	0.12	0.29
2012-2014	0.49	0.13	0.33	0.24	0.40	0.44	0.27	0.13	0.28
Reduction AAE 1995-2013	0.66	0.34	0.43	0.57	0.73	0.76	0.60	0.23	0.50




Figure 2.5: Acidity: Percentage area of habitats where acidity critical loads are exceeded and acidity AAE for the UK by deposition dataset year.

Table 2.8: Nutrient nitrogen: Percentage area of habitats where nitrogen critical loads are exceeded in the UK by deposition dataset year.

Year	Percentage habitat area with exceedance of nutrient nitrogen critical loads:												
	Acid grassland	Calcareous grassland	Dwarf shrub heath	Bog	Montane	Coniferous woodland (managed)	Broadleaved woodland (managed)	Fagus woodland (unmanaged)	Acidophilous oak (unmanaged)	Scots Pine (unmanaged)	Other unmanaged woodland	Dune grassland	Saltmarsh
1995-1997	72.6	97.5	59.1	54.2	96.7	95.4	98.4	100.0	98.9	61.1	96.5	70.6	2.0
1998-2000	61.3	95.5	49.0	45.1	95.7	90.5	97.4	100.0	97.0	38.9	95.1	44.8	1.1
1999-2001	61.4	95.5	51.1	45.0	97.1	92.8	97.8	100.0	98.1	52.3	95.5	46.9	2.1
2001-2003	63.1	95.5	47.8	44.6	89.0	90.6	97.4	100.0	96.1	49.7	95.5	41.9	1.0
2002-2004	64.3	93.9	49.8	44.9	92.6	93.0	98.1	100.0	98.2	66.5	95.7	36.1	1.1
2003-2005	64.8	93.9	50.6	45.2	90.5	92.1	98.0	100.0	98.1	67.8	95.7	33.5	1.1
2004-2006	64.8	90.6	54.5	45.9	96.6	90.2	97.5	100.0	95.6	58.0	95.5	29.3	0.8
2005-2007	64.2	89.4	54.3	54.6	96.2	91.0	97.4	100.0	95.5	52.6	95.6	31.8	0.8
2006-2008	60.0	87.7	49.5	55.4	95.5	89.4	97.1	100.0	93.8	34.2	95.5	31.1	0.8
2007-2009	56.3	89.6	43.9	47.1	82.7	86.9	96.7	100.0	89.8	30.7	95.2	29.2	0.9
2008-2010	55.7	91.2	42.7	45.6	81.0	86.1	96.7	99.9	88.5	30.5	95.1	34.7	0.9
2009-2011	61.1	92.3	45.0	45.8	82.1	88.2	97.0	99.9	91.5	32.4	95.3	37.6	0.9
2010-2012	59.7	90.4	42.2	44.8	74.4	86.5	96.8	99.9	87.7	26.2	94.7	34.0	0.9
2011-2013	60.8	87.6	41.6	43.1	71.2	86.4	96.8	100.0	88.6	24.2	95.0	29.2	0.8
2012-2014	59.8	88.7	41.2	45.2	75.7	85.1	96.5	100.0	86.6	26.0	94.9	26.5	0.7
Reduction in % area exceeded 1995-2013	13.3	8.8	17.9	9.0	21.0	10.2	1.9	0.0	12.3	35.0	1.6	44.1	1.3

Table 2.9: Nutrient nitrogen: AAE (in kg N ha⁻¹ year⁻¹) by habitat for the UK by deposition dataset year.

Year	AAE (kg N ha ⁻¹ year ⁻¹) by habitat:												
	Acid grassland	Calcareous grassland	Dwarf shrub heath	Bog	Montane	Coniferous woodland (managed)	Broadleaved woodland (managed)	Fagus woodland (unmanaged)	Acidophilous oak (unmanaged)	Scots Pine (unmanaged)	Other unmanaged woodland	Dune grassland	Saltmarsh
1995-1997	6.3	7.6	4.5	5.3	5.5	16.8	24.5	22.7	19.9	3.3	23.2	2.71	0.04
1998-2000	3.9	7.3	3.1	3.8	4.4	12.1	21.8	19.5	16.4	2.0	21.1	1.63	0.05
1999-2001	4.0	7.7	3.2	3.9	5.0	12.8	22.7	20.3	17.3	2.8	22.0	1.74	0.06
2001-2003	4.5	8.9	3.6	4.4	5.7	14.4	25.8	22.9	18.8	3.2	25.2	1.52	0.03
2002-2004	4.2	6.9	3.4	3.8	6.1	14.7	24.5	22.1	19.1	4.0	23.8	0.93	1.66
2003-2005	4.4	6.9	3.4	3.8	6.1	14.8	24.8	22.6	19.4	3.7	24.1	0.93	1.67
2004-2006	4.4	5.7	3.4	3.9	6.4	12.2	19.3	15.8	15.5	2.6	18.7	0.75	0.03
2005-2007	4.3	5.7	3.3	4.0	5.5	12.3	19.4	15.4	15.4	2.3	19.1	0.80	0.04
2006-2008	3.9	5.2	3.0	4.0	4.3	11.5	18.2	14.0	14.2	1.9	18.1	0.74	0.04
2007-2009	3.5	5.3	2.6	3.5	3.3	10.8	18.3	14.4	13.9	1.6	18.5	0.77	0.04
2008-2010	3.4	5.5	2.6	3.5	3.3	10.9	18.5	14.6	13.9	1.7	18.9	0.86	0.05
2009-2011	3.9	5.9	3.0	3.9	3.6	11.8	19.4	15.2	14.7	1.9	19.9	1.04	0.06
2010-2012	3.7	5.3	2.8	3.7	2.9	11.2	18.1	13.9	13.7	1.6	18.4	0.91	0.05
2011-2013	3.7	4.9	2.8	3.7	2.9	11.0	17.3	13.3	13.5	1.5	17.5	0.76	0.03
2012-2014	3.5	4.9	2.7	3.6	2.9	10.5	16.5	12.5	12.7	1.6	16.5	0.7	0.02
Reduction in AAE 1995-2013	2.7	2.7	1.8	1.7	2.6	6.3	8.0	10.2	7.2	1.7	6.7	2.0	0.02



Figure 2.6: Nutrient nitrogen: Percentage area of habitats where nutrient nitrogen critical loads are exceeded and nutrient nitrogen AAE (in kg N ha⁻¹ year⁻¹) in the UK by deposition dataset year.

3. Trends in exceedance of site-relevant critical loads (SRCL)

3.1 Overview of site-relevant critical loads

Site relevant critical loads (SRCL) have been applied to three types of statutory protected sites:

- Special Areas of Conservation (SACs) are protected sites designated under the EC Habitats Directive. Annexes I and II of the Directive identify the habitats and species (excluding birds) to be protected; 78 Annex I habitat types and 41 species are believed to occur in, or be native to the UK.
- Special Protected Areas (SPAs) are sites classified under the EC Birds Directive to protect rare and vulnerable birds (as listed in an Annex to the Directive) and regularly occurring migratory species.
- Sites of Special Scientific Interest (SSSIs in England, Wales and Scotland) and Areas of Special Scientific Interest (ASSIs in Northern Ireland) provide statutory protection to the UK's flora and fauna. There are additional SSSIs designated for geological or physiographic features but these are not included in the SRCL assessments.

Digital boundaries for all sites in the UK have been collated by JNCC, together with tables identifying the designated feature habitats and species associated with each site, but no digital information is currently available on the spatial area of each feature within each site. ***Therefore, for the purposes of the national SRCL work described here, it is assumed that all features recorded for a site, occur across the entire site area. To avoid double counting the area exceeding critical loads for sites with more than one designated feature, the maximum area exceeded for any feature is used when summarising results to the site and country levels (Section 3.2).***

To assign SRCL, the first step is to consider if the interest feature is potentially sensitive to acidification and/or eutrophication. Specialists within Natural England, Scottish Natural Heritage and CEH have used expert judgement to determine this (SNIFFER, 2007). For SPAs where the features are bird species, the broad habitats the birds depend upon for feeding, breeding and roosting are considered.

To assign critical loads to the habitat features of designated sites it is necessary to link the different habitat classifications used. Acidity critical loads are mapped by broad habitat and empirical critical loads of nitrogen are based on the EUNIS (European Nature Information System; Davies & Moss, 2002) habitat classification. Look-up tables developed by Davies & Moss (2002) and published in the National Biodiversity Network (NBN) Habitats Dictionary (<http://habitats.nbn.org.uk/>) and available from the JNCC website (<http://jncc.defra.gov.uk/page-1425>) enable linkages to be made between:

- Annex I habitats and EUNIS classes
- Annex I habitats and broad habitats
- EUNIS habitats and broad habitats

Using the look up tables the most appropriate EUNIS class and broad habitat class can be assigned to each interest feature. It should be noted that some sites may contain features sensitive to acidification and/or eutrophication for which no appropriate critical loads are available.

The critical loads assigned to the habitat features are based on the same methods and data as those outlined in Section 1.1 of this report. However, the national critical load maps are based on national scale data sets appropriate for national scale critical load and critical level assessments. This means they may not include all small areas of sensitive habitats or some coastal habitats; therefore some designated sites and/or feature habitats may not be included in the areas mapped nationally for critical loads. To overcome this, for SRCL a separate database of national critical loads for terrestrial habitats has been created, that provides critical loads for every 1x1km square in the UK whether the habitat is known to exist there or not. The appropriate SRCL can then be extracted for terrestrial habitat features of each designated site; the SRCL does not include any acidity critical loads for freshwater habitats. For further information refer to the “Methods” report (Hall et al, 2015).

For nutrient nitrogen the empirical critical loads approach is applied to designated feature habitats sensitive to nitrogen. The critical load value applied to each habitat are the “Recommended” values agreed by habitat specialists for Article 17 reporting (for more information refer to <http://www.apis.ac.uk/indicative-critical-load-values> and to Hall et al, 2015).

3.2 Overview of SRCL exceedance metrics

Exceedances are calculated separately for SACs, SPAs and SSSIs, for all site features that critical loads and deposition data can be assigned to (Hall et al, 2015). Metrics are calculated by:

(a) Feature (within each site):

- Exceedance
- Exceeded area[#]
- Accumulated Exceedance (AE)(i.e. exceedance * exceeded area)
- Average Accumulated Exceedance (AAE)(i.e. AE / total site area)

(b) Site:

- Total number of features with SRCL
- Number and percentage of features with exceedance of SRCL.
- Maximum area exceeded^{##} for any feature within a site
- Maximum AE for any feature within a site
- Maximum AAE for any feature within a site

(c) Country:

- Total number of sites
- Total number and percentage of sites with SRCL for one or more features
- Total number of features with SRCL
- Total number and percentage of sites with exceedance of SRCL for one or more features
- Total number and percentage of features with exceedance of SRCL
- Total area of all sites
- Total area of all sites with SRCL
- Maximum exceeded area^{###}
- Maximum AE calculated as the sum of the maximum AE for all sites
- Maximum AAE; calculated from the country maximum AE and total area of all sites (with SRCL) within a country.

Feature exceeded area: If the critical load is exceeded and the deposition values are constant across the whole site, the exceeded area equals the site area; if the deposition values vary across the site (e.g. as a result of the site crossing the boundaries between different 5x5km grid squares with different deposition values), then the exceeded area will be the sum of the 1x1 km portions of the site where the deposition exceeds the critical load.

Site maximum exceeded area: is set to the maximum exceeded area for any feature within a site.

Country maximum exceeded area: is calculated as the sum of the site maximum exceeded areas for all sites within a country.

The sections below summarise the results by country, based on the CBED deposition (Section 1.2) for 1995-97 to 2012-14. Note that the summary statistics may present the “worst” case, as they include all sites where at least one feature is exceeded (other features within sites may have smaller or no exceedance), and the AAE results are based on the maximum exceedance of any feature within a site.

3.2.1 Acidity results

The trends in acidity critical load exceedances are summarised in Tables 3.1-3.3 and present the percentage of sites (with SRCL) by country, where the SRCL is exceeded for one or more features, together with the maximum AAE. For all site types (SACs, SPAs and SSSIs) the largest reductions in the percentage of sites with critical load exceedance between 1995 and 2014 are for Scotland (26-35% reduction), and the AAE has at least halved in all countries over this time period. These trends reflect the changing patterns of acid deposition over this time period. At the UK level the latest results (based on CBED deposition for 2012-14) show (i) for SACs, 76% of sites have exceedance of the acidity critical loads for one or more features (down 15% from 1995), and a maximum AAE of 0.66 keq ha⁻¹ year⁻¹ (down by 0.85 keq ha⁻¹ year⁻¹ from 1995); (ii) for SPAs, 70.3% of sites have exceedance of the acidity critical loads for one or more features (down 24% from 1995), and a maximum AAE of 0.46 keq ha⁻¹ year⁻¹ (down by 0.65 keq ha⁻¹ year⁻¹ from 1995); (iii) for SSSIs, 61.4% of sites have exceedance of the acidity critical loads for one or more features (down 16.1% from 1995), and a maximum AAE of 0.48 keq ha⁻¹ year⁻¹ (down by 0.7 keq ha⁻¹ year⁻¹ from 1995).

Maps of the maximum AAE per site (Figure 3.1) based on the latest CBED deposition (2012-14) show the highest exceedances in northern England and parts of Wales, with the lowest exceedances across Scotland. Some SACs and SPAs in the far north of Scotland, and some small SSSIs across parts of Scotland and southern England have no exceedance for any feature.

Table 3.1: Trends in acidity exceedances for SACs; percentage of sites (with SRCL) with exceedance of SRCL for at least one feature, and maximum AAE of all sites/features.

Parameter	Country						
	England	Wales	Scotland	NI	Eng/Wales*	Eng/Scot*	UK
Number of sites	231	85	236	54	7	3	616
Number of sites with SRCL for at least one feature	180	71	182	47	6	1	487
(a) % of sites with SRCL with exceedance of SRCL for at least one feature, (b)[maximum AAE keq ha ⁻¹ year ⁻¹], by deposition data for:							
1995-1997	85.0 [2.36]	97.2 [1.87]	92.3 [0.66]	97.9 [1.32]	100.0	100.0	91.0 [1.51]
1998-2000	82.2 [1.80]	97.2 [1.29]	83.5 [0.42]	95.7 [0.76]	100.0	100.0	86.4 [1.10]
1999-2001	81.7 [1.83]	97.2 [1.31]	83.5 [0.44]	95.7 [0.78]	100.0	100.0	86.2 [1.12]
2001-2003	81.1 [1.89]	94.4 [1.31]	75.3 [0.41]	95.7 [0.87]	100.0	100.0	82.5 [1.13]
2002-2004	82.8 [1.77]	95.8 [1.27]	78.0 [0.43]	95.7 [0.77]	100.0	100.0	84.4 [1.09]
2003-2005	82.8 [1.75]	95.8 [1.33]	76.4 [0.42]	95.7 [0.71]	100.0	100.0	83.8 [1.08]
2004-2006	79.4 [1.50]	95.8 [1.08]	79.7 [0.42]	95.7 [0.70]	100.0	100.0	83.8 [0.95]
2005-2007	79.4 [1.45]	95.8 [1.05]	79.7 [0.38]	95.7 [0.73]	100.0	100.0	83.8 [0.91]
2006-2008	77.2 [1.35]	95.8 [0.90]	75.8 [0.31]	95.7 [0.71]	100.0	100.0	81.5 [0.82]
2007-2009	76.7 [1.21]	95.8 [0.82]	69.2 [0.22]	95.7 [0.72]	100.0	100.0	78.9 [0.71]
2008-2010	75.6 [1.16]	95.8 [0.77]	67.6 [0.22]	95.7 [0.75]	100.0	100.0	77.8 [0.68]
2009-2011	76.1 [1.20]	95.8 [0.75]	70.3 [0.23]	95.7 [0.79]	100.0	100.0	79.1 [0.71]
2010-2012	76.1 [1.17]	93.0 [0.75]	68.1 [0.21]	93.6 [0.72]	100.0	100.0	77.6 [0.68]
2011-2013	75.0 [1.18]	93.0 [0.75]	68.1 [0.19]	95.7 [0.72]	100.0	100.0	77.4 [0.67]
2012-2014	74.4 [1.14]	94.4 [0.79]	64.3 [0.19]	95.7 [0.63]	100.0	100.0	76.0 [0.66]
Reduction in % sites with exceedance 1995-2014 [Reduction in maximum AAE keq ha ⁻¹ year ⁻¹ 1995-2014]	10.6 [1.22]	2.8 [1.07]	28.0 [0.47]	2.1 [0.69]	0.0	0.0	15.0 [0.85]

* Some sites cross the England/Wales or England/Scotland border and have been assigned to these border areas. However, in calculating AAE each 1x1 km square (or part thereof) within each site has been assigned to a single country, so AAE results are calculated for individual countries only.

Table 3.2: Trends in acidity exceedances for SPAs; percentage of sites (with SRCL) with exceedance of SRCL for at least one feature, and maximum AAE of all sites/features.

Parameter	Country						
	England	Wales	Scotland	NI	Eng/Wales*	Eng/Scot*	UK
Number of sites	78	17	145	14	3	0	257
Number of sites with SRCL for at least one feature	63	13	86	10	3	0	175
(a) % of sites with SRCL with exceedance of SRCL for at least one feature, (b)[maximum AAE keq ha ⁻¹ year ⁻¹], by deposition data for:							
1995-1997	98.4 [1.73]	100.0 [1.85]	89.5 [0.45]	100.0 [1.09]	100.0	-	94.3 [1.11]
1998-2000	98.4 [1.41]	100.0 [1.14]	72.1 [0.24]	100.0 [0.43]	100.0	-	85.7 [0.81]
1999-2001	98.4 [1.39]	100.0 [1.21]	73.3 [0.24]	90.0 [0.41]	100.0	-	85.7 [0.80]
2001-2003	98.4 [1.38]	84.6 [1.24]	62.8 [0.23]	70.0 [0.55]	100.0	-	78.3 [0.80]
2002-2004	96.8 [1.24]	92.3 [1.25]	66.3 [0.23]	80.0 [0.32]	100.0	-	80.6 [0.73]
2003-2005	96.8 [1.21]	92.3 [1.32]	65.1 [0.19]	80.0 [0.26]	100.0	-	80.0 [0.70]
2004-2006	88.9 [1.08]	92.3 [1.01]	62.8 [0.19]	90.0 [0.27]	100.0	-	76.6 [0.63]
2005-2007	90.5 [1.04]	92.3 [0.98]	66.3 [0.19]	80.0 [0.25]	100.0	-	78.3 [0.61]
2006-2008	90.5 [0.99]	92.3 [0.77]	64.0 [0.16]	80.0 [0.25]	100.0	-	77.1 [0.55]
2007-2009	90.5 [0.91]	92.3 [0.66]	55.8 [0.10]	80.0 [0.23]	100.0	-	73.1 [0.48]
2008-2010	90.5 [0.87]	92.3 [0.62]	54.7 [0.09]	80.0 [0.25]	100.0	-	72.6 [0.47]
2009-2011	88.9 [0.90]	92.3 [0.59]	58.1 [0.11]	80.0 [0.32]	100.0	-	73.7 [0.48]
2010-2012	88.9 [0.89]	84.6 [0.59]	58.1 [0.09]	80.0 [0.27]	100.0	-	73.1 [0.47]
2011-2013	87.3 [0.86]	84.6 [0.59]	54.7 [0.08]	90.0 [0.28]	100.0	-	71.4 [0.45]
2012-2014	87.3 [0.88]	84.6 [0.65]	54.7 [0.08]	70.0 [0.17]	100.0	-	70.3 [0.46]
Reduction in % sites with exceedance [Reduction in maximum AAE keq ha ⁻¹ year ⁻¹ 1995-2014]	11.1 [0.85]	15.4 [1.20]	34.9 [0.37]	30.0 [0.92]	0.0	-	24.0 [0.65]

* Some sites cross the England/Wales or England/Scotland border and have been assigned to these border areas. However, in calculating AAE each 1x1 km square (or part thereof) within each site has been assigned to a single country, so AAE results are calculated for individual countries only.

Table 3.3: Trends in acidity exceedances for SSSIs; percentage of sites (with SRCL) with exceedance of SRCL for at least one feature, and maximum AAE of all sites/features.

Parameter	Country						
	England	Wales	Scotland	NI	Eng/Wales*	Eng/Scot*	UK
Number of sites	4115	1018	1452	291	0	0	6876
Number of sites with SRCL for at least one feature	2924	676	905	178	0	0	4683
(a) % of sites with SRCL with exceedance of SRCL for at least one feature, (b)[maximum AAE keq ha ⁻¹ year ⁻¹], by deposition data for:							
1995-1997	71.8 [1.66]	91.4 [1.63]	85.1 [0.60]	82.0 [1.18]	-	-	77.6 [1.18]
1998-2000	68.0 [1.24]	80.9 [1.11]	76.1 [0.37]	75.3 [0.65]	-	-	71.7 [0.84]
1999-2001	68.2 [1.24]	81.2 [1.09]	75.8 [0.36]	75.3 [0.66]	-	-	71.8 [0.83]
2001-2003	67.9 [1.27]	79.6 [1.07]	69.7 [0.32]	75.3 [0.77]	-	-	70.2 [0.82]
2002-2004	66.5 [1.16]	81.8 [1.04]	70.1 [0.33]	77.0 [0.65]	-	-	69.8 [0.78]
2003-2005	66.3 [1.15]	82.0 [1.10]	69.1 [0.32]	75.8 [0.59]	-	-	69.5 [0.77]
2004-2006	62.7 [0.99]	79.1 [0.95]	69.8 [0.33]	74.7 [0.58]	-	-	66.9 [0.69]
2005-2007	62.3 [0.95]	79.4 [0.93]	70.1 [0.31]	75.8 [0.60]	-	-	66.8 [0.66]
2006-2008	60.7 [0.88]	78.6 [0.80]	67.2 [0.25]	75.3 [0.59]	-	-	65.1 [0.59]
2007-2009	60.2 [0.79]	77.5 [0.72]	62.9 [0.17]	75.3 [0.59]	-	-	63.8 [0.51]
2008-2010	60.0 [0.76]	76.0 [0.67]	61.2 [0.16]	75.8 [0.61]	-	-	63.1 [0.49]
2009-2011	60.1 [0.79]	76.0 [0.65]	63.2 [0.19]	78.1 [0.65]	-	-	63.7 [0.51]
2010-2012	59.3 [0.77]	74.7 [0.65]	61.8 [0.17]	76.4 [0.60]	-	-	62.7 [0.50]
2011-2013	58.2 [0.77]	74.9 [0.66]	59.1 [0.16]	77.0 [0.60]	-	-	61.5 [0.49]
2012-2014	58.4 [0.75]	75.0 [0.70]	58.8 [0.16]	73.0 [0.51]	-	-	61.4 [0.48]
Reduction in % sites with exceedance [Reduction in maximum AAE keq ha ⁻¹ year ⁻¹ 1995-2014]	13.3 [0.91]	16.4 [0.93]	26.3 [0.43]	9.0 [0.67]	-	-	16.1 [0.70]

* Some SACs and SPAs cross the England/Wales or England/Scotland border and have been assigned to these border areas; all SSSIs have been assigned to a single country only.




Figure 3.1: Average Accumulated Exceedance (AAE) of acidity critical loads by CBED deposition for 2012-14; maps show the maximum AAE for any feature within each site (other features may have lower or no exceedance).

3.2.2 Nutrient nitrogen results

The trends in nutrient nitrogen critical load exceedances from 1995 to 2014 are summarised in Tables 3.4-3.6. The reductions in the percentage of sites with exceedance of nutrient nitrogen critical loads for one or more features, and reductions in AAE, are smaller than the reductions seen for acidity, reflecting the smaller decreases in nitrogen deposition over time. Reductions vary by country for the different site types; the largest reductions in the percentage of sites with exceedance is greatest in Scotland for SACs (6.5% reduction) and SSSIs (11% reduction), and in Wales (21% reduction) for SPAs, though there are fewer SPAs in Wales compared to Scotland and England. The largest reductions in AAE between 1995 and 2014 are for Scotland (37-46%) and Wales (36-41%).

At the UK level the latest results (based on CBED deposition for 2012-14) show (i) for SACs, 90.1% of sites have exceedance of the nutrient nitrogen critical loads for one or more features (down 4.9% from 1995), and a maximum AAE of 9.1 kg N ha⁻¹ year⁻¹ (down by 5.0 kg N ha⁻¹ year⁻¹ from 1995); (ii) for SPAs, 73.3% of sites have exceedance of the nutrient nitrogen critical loads for one or more features (down 10.7% from 1995), and a maximum AAE of 8.7 kg N ha⁻¹ year⁻¹ (down by 4.6 kg N ha⁻¹ year⁻¹ from 1995); (iii) for SSSIs, 88.1% of sites have exceedance of the nutrient nitrogen critical loads for one or more features (down 6.8% from 1995), and a maximum AAE of 9.7 kg N ha⁻¹ year⁻¹ (down by 5.2 kg N ha⁻¹ year⁻¹ from 1995).

Maps of the maximum AAE per site (Figure 3.2) based on the latest CBED deposition (2012-14) show very few sites with no exceedance of any feature. Exceedances are widespread across all countries with generally lower exceedances in Scotland. The maximum AAE is above 7 kg N ha⁻¹ year⁻¹ for the majority of sites, with many sites having maximum AAE up to 28 kg N ha⁻¹ year⁻¹, and a few sites in central England with maximum AAE above this.

Table 3.4: Trends in nutrient nitrogen exceedances for SACs; percentage of sites (with SRCL) with exceedance of SRCL for at least one feature, and maximum AAE of all sites/features.

Parameter	Country						
	England	Wales	Scotland	NI	Eng/Wales*	Eng/Scot*	UK
Number of sites	231	85	236	54	7	3	616
Number of sites with SRCL for at least one feature	197	79	201	50	7	2	536
(a) % of sites with SRCL with exceedance of SRCL for at least one feature, (b)[maximum AAE kg N ha ⁻¹ year ⁻¹], by deposition data for:							
1995-1997	98.5 [20.5]	98.7 [14.1]	89.6 [7.3]	98.0 [14.4]	100.0	50.0	95.0 [14.1]
1998-2000	97.0 [17.4]	96.2 [10.3]	85.1 [5.9]	96.0 [9.8]	100.0	50.0	92.2 [11.4]
1999-2001	97.0 [18.3]	96.2 [10.9]	85.6 [6.3]	96.0 [10.3]	100.0	50.0	92.4 [12.1]
2001-2003	98.0 [19.8]	94.9 [11.9]	84.6 [6.3]	98.0 [12.7]	100.0	50.0	92.4 [12.9]
2002-2004	97.5 [18.4]	93.7 [11.7]	85.6 [6.6]	98.0 [11.3]	100.0	50.0	92.4 [12.4]
2003-2005	97.5 [18.6]	96.2 [12.2]	85.6 [6.4]	98.0 [10.9]	100.0	50.0	92.7 [12.5]
2004-2006	95.9 [15.8]	94.9 [9.7]	84.6 [6.2]	98.0 [11.0]	100.0	50.0	91.6 [10.9]
2005-2007	94.9 [15.7]	94.9 [9.7]	86.1 [6.5]	98.0 [11.9]	100.0	50.0	91.8 [11.0]
2006-2008	94.4 [15.0]	93.7 [8.8]	86.6 [6.1]	98.0 [11.8]	100.0	50.0	91.6 [10.4]
2007-2009	94.9 [14.1]	93.7 [8.5]	83.1 [5.1]	98.0 [12.3]	100.0	50.0	90.5 [9.6]
2008-2010	95.4 [14.0]	93.7 [8.4]	82.6 [4.8]	98.0 [12.7]	100.0	50.0	90.5 [9.4]
2009-2011	95.9 [14.5]	93.7 [8.4]	84.1 [4.9]	98.0 [13.1]	100.0	50.0	91.2 [9.7]
2010-2012	95.4 [14.0]	93.7 [8.2]	83.1 [4.6]	98.0 [12.2]	100.0	50.0	90.7 [9.3]
2011-2013	93.9 [13.9]	93.7 [8.2]	82.6 [4.4]	98.0 [12.2]	100.0	50.0	89.9 [9.2]
2012-2014	94.4 [13.5]	93.7 [8.3]	83.1 [4.6]	96.0 [11.1]	100.0	50.0	90.1 [9.1]
Reduction in % sites with exceedance 1995-2014 [Reduction in maximum AAE kg N ha ⁻¹ year ⁻¹ 1995-2014]	4.1 [7.0]	5.1 [5.8]	6.5 [2.7]	2.0 [3.3]	0.0	0.0	4.9 [5.0]

* Some sites cross the England/Wales or England/Scotland border and have been assigned to these border areas. However, in calculating AAE each 1x1 km square (or part thereof) within each site has been assigned to a single country, so AAE results are calculated for individual countries only.

Table 3.5: Trends in nutrient nitrogen exceedances for SPAs; percentage of sites (with SRCL) with exceedance of SRCL for at least one feature, and maximum AAE of all sites/features.

Parameter	Country						
	England	Wales	Scotland	NI	Eng/Wales*	Eng/Scot*	UK
Number of sites	78	17	145	14	3	0	257
Number of sites with SRCL for at least one feature	72	14	124	12	3	0	225
(a) % of sites with SRCL with exceedance of SRCL for at least one feature, (b)[maximum AAE kg N ha ⁻¹ year ⁻¹], by deposition data for:							
1995-1997	97.2 [18.7]	100.0 [18.5]	74.2 [6.5]	83.3 [14.9]	100.0	-	84.0 [13.3]
1998-2000	94.4 [17.4]	100.0 [13.4]	69.4 [4.9]	83.3 [8.9]	100.0	-	80.4 [11.2]
1999-2001	97.2 [17.8]	100.0 [14.4]	69.4 [5.2]	83.3 [9.1]	100.0	-	81.3 [11.7]
2001-2003	95.8 [18.7]	85.7 [15.9]	62.9 [5.2]	83.3 [13.6]	100.0	-	76.4 [12.4]
2002-2004	93.1 [17.1]	100.0 [16.2]	66.9 [5.2]	83.3 [11.1]	100.0	-	78.7 [11.6]
2003-2005	93.1 [17.2]	100.0 [17.0]	65.3 [4.9]	83.3 [10.7]	100.0	-	77.8 [11.5]
2004-2006	93.1 [14.3]	92.9 [13.7]	66.9 [4.8]	83.3 [10.1]	100.0	-	78.2 [10.0]
2005-2007	90.3 [14.2]	100.0 [13.7]	68.5 [5.2]	83.3 [11.1]	100.0	-	78.7 [10.1]
2006-2008	88.9 [13.6]	100.0 [12.4]	69.4 [4.9]	83.3 [11.1]	66.7	-	78.2 [9.7]
2007-2009	91.7 [13.3]	100.0 [11.9]	65.3 [4.0]	83.3 [11.6]	100.0	-	77.3 [9.1]
2008-2010	91.7 [13.4]	100.0 [11.6]	62.9 [3.8]	83.3 [12.0]	100.0	-	76.0 [9.1]
2009-2011	93.1 [13.8]	100.0 [11.3]	68.5 [4.0]	91.7 [13.3]	100.0	-	80.0 [9.4]
2010-2012	90.3 [13.3]	92.9 [11.2]	62.9 [3.7]	83.3 [12.2]	100.0	-	75.1 [9.0]
2011-2013	90.3 [12.9]	92.9 [11.3]	64.5 [3.4]	83.3 [11.8]	100.0	-	76.0 [8.7]
2012-2014	88.9 [13.0]	78.6 [11.8]	62.1 [3.5]	83.3 [9.9]	100.0	-	73.3 [8.7]
Reduction in % sites with exceedance 1995-2014 [Reduction in maximum AAE kg N ha ⁻¹ year ⁻¹ 1995-2014]	8.3 [5.7]	21.4 [6.6]	12.1 [3.0]	0 [5.0]	0	-	10.7 [4.6]

* Some sites cross the England/Wales or England/Scotland border and have been assigned to these border areas. However, in calculating AAE each 1x1 km square (or part thereof) within each site has been assigned to a single country, so AAE results are calculated for individual countries only.

Table 3.6: Trends in nutrient nitrogen exceedances for SSSIs; percentage of sites (with SRCL) with exceedance of SRCL for at least one feature, and maximum AAE of all sites/features.

Parameter	Country						
	England	Wales	Scotland	NI	Eng/Wales*	Eng/Scot*	UK
Number of sites	4115	1018	1452	291	0	0	6876
Number of sites with SRCL for at least one feature	2954	686	938	188	0	0	4766
(a) % of sites with SRCL with exceedance of SRCL for at least one feature, (b)[maximum AAE kg N ha ⁻¹ year ⁻¹], by deposition data for:							
1995-1997	95.2 [20.9]	100.0 [20.7]	90.1 [7.0]	94.1 [16.3]	-	-	94.9 [14.9]
1998-2000	94.2 [18.1]	98.8 [15.3]	85.0 [5.3]	86.2 [10.9]	-	-	92.7 [12.2]
1999-2001	94.8 [18.8]	98.8 [16.0]	85.9 [5.7]	87.8 [11.3]	-	-	93.3 [12.8]
2001-2003	96.4 [20.4]	97.4 [17.6]	82.9 [5.6]	87.8 [15.2]	-	-	93.6 [13.8]
2002-2004	92.6 [18.8]	98.8 [17.8]	84.3 [5.9]	88.8 [14.0]	-	-	91.7 [13.1]
2003-2005	93.2 [19.1]	98.8 [18.7]	83.6 [5.7]	88.8 [13.6]	-	-	91.9 [13.3]
2004-2006	89.4 [16.2]	98.7 [15.7]	83.2 [5.6]	88.3 [12.8]	-	-	89.4 [11.6]
2005-2007	89.9 [16.1]	99.0 [15.7]	84.6 [5.8]	89.4 [14.1]	-	-	90.2 [11.7]
2006-2008	88.6 [15.3]	98.8 [14.2]	83.7 [5.3]	92.0 [14.0]	-	-	89.2 [11.0]
2007-2009	89.5 [14.6]	98.5 [13.7]	81.2 [4.3]	93.1 [14.7]	-	-	89.3 [10.3]
2008-2010	89.9 [14.6]	98.4 [13.4]	80.3 [4.1]	93.1 [15.1]	-	-	89.4 [10.1]
2009-2011	90.0 [15.2]	98.5 [13.4]	81.1 [4.4]	93.1 [16.2]	-	-	89.8 [10.6]
2010-2012	89.1 [14.6]	98.0 [13.0]	80.1 [4.1]	92.6 [14.8]	-	-	88.8 [10.1]
2011-2013	87.7 [14.4]	98.1 [13.2]	80.2 [4.0]	93.1 [14.5]	-	-	87.9 [9.9]
2012-2014	88.5 [14.0]	98.3 [13.4]	79.1 [4.0]	89.9 [12.6]	-	-	88.1 [9.7]
Reduction in % sites with exceedance 1995-2014 [Reduction in maximum AAE kg N ha ⁻¹ year ⁻¹ 1995-2014]	6.8 [6.9]	1.7 [7.4]	11.0 [3.0]	4.3 [3.7]	-	-	6.8 [5.2]

* Some SACs and SPAs cross the England/Wales or England/Scotland border and have been assigned to these border areas; all SSSIs have been assigned to a single country only.




Figure 3.2: Average Accumulated Exceedance (AAE) of nutrient nitrogen critical loads by CBED deposition for 2012-14; maps show the maximum AAE for any feature within each site (other features may have lower or no exceedance).

References

- Bobbink, R. & Hettelingh, J.P. (eds) 2011. Review and revision of empirical critical loads and dose-response relationships. Coordination Centre for Effects, National Institute for Public Health and the Environment (RIVM). <http://wge-cce.org>
- Calver, L. 2003. A suggested improved method for the quantification of critical loads of acidity for peat soils. PhD Thesis, University of York.
- Calver, L.J., Cresser, M.S. & Smart, R.P. 2004. Tolerance of *Calluna vulgaris* and peatland plant communities to sulphuric acid deposition. *Chemistry and Ecology*, 20, 309-320.
- Davies, C.E. & Moss, D. 2002. EUNIS Habitat Classification. 2001 Work Programme, Final Report to the European Environment Agency European Topic Centre on Nature Protection and Biodiversity. Centre for Ecology and Hydrology, February 2002.
- Fowler, D., Cape, J.N., Leith, I.D., Choularton, T.W., Gay, M.J. & Jones, A. 1988. The influence of altitude on rainfall composition at Great Dun Fell. *Atmospheric Environment*, 22, 1355-1362.
- Fuller, R.M., Smith, G.M., Sanderson, J.M., Hill, R.A. & Thomson, A.G. 2002a. The UK Land Cover Map 2000: construction of a parcel based vector map from satellite images. *Cartographic Journal*, 39, 115-25.
- Fuller, R.M., Smith, G.M., Sanderson, J.M., Hill, R.A., Thomson, A.G., Cox, R., Brown, N.J. & Gerard, F. 2002b. Countryside Survey 2000 Module 7: Land Cover Map 2000. Final Report, CSLCM Final CEH report to Defra.
- Gammack, S.M., Smith, C.M.S. and Cresser, M.S. 1995. The approach used for mapping critical loads for ombrotrophic peats in Great Britain, *Proceedings of a Conference on: Acid Rain and its Impact: The Critical Loads Debate*, R.W. Battarbee (Ed.), 180-183, Ensis Publishing, London.
- Hall, J., Curtis, C., Dore, T. & Smith, R. 2015. Methods for the calculation of critical loads and their exceedances in the UK. Report to Defra under contract AQ0826. CEH Bangor. <http://www.cldm.ceh.ac.uk>
- Henriksen, A. & Posch, M. 2001. Steady-state models for calculating critical loads of acidity for surface waters. *Water, Air and Soil Pollution: Focus* 1, 375-398.
- Hettelingh, J.-P., Posch, M., de Smet, P.A.M. & Downing, R.J. 1995. The use of critical loads in emission reduction agreements in Europe. *Water, Air and Soil Pollution*, 85, 2381-2388.
- Hornung, M., Bull, K., Cresser, M., Hall, J., Langan, S., Loveland, P. and Smith, C. 1995c. An empirical map of critical loads for soils in Great Britain, *Environmental Pollution*, 90, 301-310.
- Nilsson, J. & Grennfelt, P. 1988, Critical loads for sulphur and nitrogen. Report 1988:15. UNECE/Nordic Council of Ministers, Copenhagen, Denmark.
- Posch, M., de Vries, W. & Hettelingh, J.-P. 1995. Critical loads of sulphur and nitrogen. In: Posch, M., de Smet, P.A.M., Hettelingh, J.-P. & Downing, R.J. (Eds.), *Calculation and Mapping of Critical Thresholds in Europe: Status Report 1995*. Coordination Centre for Effects, RIVM, Bilthoven, Netherlands. pp 31-41. <http://wge-cce.org/>

Posch, M., de Smet, P.A.M., Hettelingh, J.-P. & Downing, R.J. (Eds.) 1999. Calculation and Mapping of Critical Thresholds in Europe: Status Report 1999. Coordination Centre for Effects, RIVM, Bilthoven, Netherlands. <http://wge-cce.org/>

Posch, M. & Hettelingh, J.-P. 1997. Remarks on critical load calculations. In: Posch, M., de Smet, P.A.M., Hettelingh, J.-P. & Downing, R.J. (Eds.), Calculation and Mapping of Critical Thresholds in Europe: Status Report 1997. Coordination Centre for Effects, RIVM, Bilthoven, Netherlands. pp 25-28. <http://wge-cce.org/>

Posch, M., de Smet, P.A.M. & Hettelingh, J.-P. 1999. Critical loads and their exceedances in Europe: an overview. In: Posch, M., de Smet, P.A.M., Hettelingh, J.-P. & Downing, R.J. (Eds.), Calculation and Mapping of Critical Thresholds in Europe: Status Report 1999. Coordination Centre for Effects, RIVM, Bilthoven, Netherlands. pp 3-11. <http://wge-cce.org/>

RoTAP. 2012. Review of Transboundary Air Pollution: Acidification, Eutrophication, Ground Level Ozone and Heavy Metals in the UK. Contract Report to the Department for Environment, Food and Rural Affairs. Centre for Ecology and Hydrology. www.rotap.ceh.ac.uk

Skiba, U. & Cresser, M. 1989. Prediction of long-term effects of rainwater acidity on peat and associated drainage water chemistry in upland areas. *Water Research*, 23, 1477-1482.

Smith, C.M.S., Cresser, M.S. and Mitchell, R.D.J. 1992. Sensitivity to acid deposition of dystrophic peat in Great Britain, *Ambio*, **22**, 22-26.

Smith, R.I., Fowler, D., Sutton, M.A., Flechard, C. & Coyle, M. 2000. Regional estimation of pollutant gas deposition in the UK: model description, sensitivity analyses and outputs. *Atmospheric Environment*, 34, 3757-3777.

Sverdrup, H., De Vries, W. & Henriksen, A. 1990. Mapping critical loads. Guidance to criteria, methods and examples for mapping critical loads and areas where they have been exceeded. Annex to the UNECE Task Force on Mapping manual on methodologies and criteria for mapping critical levels/loads and geographical areas where they are exceeded. Report 1990: 14, Nord 1990: 98. Copenhagen: Nordic Council of Ministers.

Sverdrup, H. & De Vries, 1994. Calculating critical loads for acidity with the simple mass balance method. *Water, Air and Soil Pollution*, 72, 143-162.